

# Influence of organic manures and bio-fertilisers on yield attributes and economics of *kharif* onion (*Allium cepa* L.) in semi-arid region

A K Meena\*, R Paliwal, K K Meena

SKN College of Agriculture, Jobner, Rajasthan 303329.

\*Corresponding author's email: ashokageco@gmail.com

(Received: 16.09.2015; Accepted: 23.02.2016)

Onion (*Allium cepa* L.) is a bulbous biennial herb of family Alliaceae. It is commonly called as "Queen of kitchen" for its unique usage throughout the year in the form of salads, condiments or for cooking with other vegetables. The pungency in onion is due to sulphur compound "ally propyl disulphide" in the volatile oil and the outer skin colour is due to the presence of "querctin" (Nadkarni, 1954). Onion bulb is rich in minerals like phosphorus (50mg/100g), iron (0.7mg/100g), calcium (18mg/100g), carbohydrates (11.0g/100g), protein (1.2g/100g), vitamins 'C' (11mg/100g), fibers (0.6g/100g) and nicotinic acid (0.4mg/100g) (Aykroyd, 1963). The productivity of onion in India is very low (16 t ha<sup>-1</sup>) in comparison to other countries. Thus there is ample scope for increasing production through fertilizers, especially that of organic manures and bio-fertilisers in light textured soil. Production of onion in *kharif* season is more important to have continuous supply of onion round the year. Organic manures stimulates the production of polysaccharides and other compounds that favours aggregation of fine soil particles, thereby promoting good structure, improved tilth, aeration, moisture movement and retention (Bose *et al.*, 2001). Bio-fertiliser inoculation like *Azospirillum*, PSB helps the plants to attain better vegetative growth and increases yield by 10-30 percent (Mohondas, 1999 and Tilak and Annapurna, 1993). The present investigation was taken up to study effect of organic manures and bio-fertilisers on *kharif* onion in semi arid region.

The experiment was conducted during 2011-12 at Horticulture farm, S.K.N. College of Agriculture, Jobner (Jaipur) during *rabi* season. The soil of experimental field was alkaline loamy sand in texture at pH 8.1, poor in organic carbon (0.135 %), available N (134.70 kg/ha), P (16.85 kg/ha), K (151.65 kg/ha) and Zn (0.42 mg/kg soil). The experiment was laid out in randomized block design (RBD) with eighteen treatment combinations including six levels of organic manures (Control, FYM @ 10 t ha<sup>-1</sup>, vermicompost @ 5 t ha<sup>-1</sup>, poultry manure @ 5 t ha<sup>-1</sup>, FYM @ 5 t ha<sup>-1</sup> + vermicompost @ 2.5 t ha<sup>-1</sup>, FYM @ 5 t ha<sup>-1</sup> + poultry manure @ 2.5 t ha<sup>-1</sup>) and three bio-fertilizer treatments (without inoculation, *Azospirillum*, *Azospirillum* + PSB) with 3 replications. Randomization of the treatments was done with the help of random number table as advocated by Fisher (1950). The plot

size was 2.1m X 2.4 m with 35 cm × 15 cm spacing between rows and plants. Organic manures were spread in the beds uniformly before transplanting of seedling. *Azospirillum* and PSB were applied as 100g per acre culture dissolved of water and dipping the bulb of the onion in solution for 10-20 minutes before sowing and dried in shade (Paul *et al.*, 1971). Yield attributes were calculated using standard methods. For calculating economics, the average treatment yield along with prevailing market rates of the produce and cost of inputs were used. The 'F-test' and critical difference (CD) calculated to test significance of difference among the treatments, wherever the results were significant.

## Effect of organic manures on yield and yield attributes

The combined application of FYM @ 5 t ha<sup>-1</sup> + poultry manure @ 2.5 t ha<sup>-1</sup> registered significantly superior increase in yield attributes over other doses and control. The increased yield and yield attributes with combined application might be due to rapid availability and utilization of nutrients, by creation of favourable soil environment, for various metabolic plant processes for carbohydrate production, which undergo hydrolysis and get distributed to reproductive parts which ultimately helped in increasing yield and may also be due to the increased rate of release of macro and micronutrients during the course of microbial decomposition (Choudhary *et al.*, 2003 Narayanamma *et al.*, 2004 and Blay *et al.*, 2002).

## Effect of bio-fertilisers on yield and yield attributes

Combined inoculation with *Azospirillum* + PSB significantly increased the neck thickness, diameter of bulb, fresh weight of bulb and bulb yield over other treatments (Table 1). The increased availability of nitrogen due to *Azospirillum* coupled with phosphorus due to PSB might have increased the yield attributes and ultimately the yield. These findings corroborate the results of Mehtha *et al.*, 1995; Subbian, 1994 and Pareek *et al.*, 1996.

## Effect of organic manures and bio-fertilisers on economics

The highest B: C ratio (3.34:1) was obtained with application of FYM @ 5 t ha<sup>-1</sup> + poultry manure @ 2.5 t ha<sup>-1</sup>, and *Azospirillum* + PSB (2.93: 1). This might be due to the



Table 1. Influence of organic manures and bio-fertilisers on growth and quality attributes of kharif onion (*Allium cepa* L.)

Treatments	Neck thickness of bulb (cm)	Diameter of bulb (cm)	Fresh weight of bulb (g)	Total bulb yield (q ha <sup>-1</sup> )	Net returns	B: C ratio
Organic manures						
Control	0.95	3.68	79.78	136.30	111210	2.12
FYM 10 t ha <sup>-1</sup>	1.07	4.20	88.44	155.83	132650	2.44
Vermicompost 5 t ha <sup>-1</sup>	1.08	4.55	92.50	160.61	130383	2.09
Poultry manure 5 t ha <sup>-1</sup>	1.18	5.06	101.33	179.61	159683	2.86
FYM 5 t ha <sup>-1</sup> + V.C. 2.5 t ha <sup>-1</sup>	1.20	5.37	103.56	187.61	166783	2.86
FYM 5 t ha <sup>-1</sup> + P.M. 2.5 t ha <sup>-1</sup>	1.28	5.54	109.68	199.50	184300	3.34
SEm±	0.03	0.15	2.74	6.30	7555	0.14
CD (p=0.05)	0.010	0.43	7.87	18.10	21714	0.40
Bio-fertilisers						
Control	1.02	4.12	89.64	155.22	129975	2.31
<i>Azospirillum</i>	1.14	4.68	95.34	169.70	147228	2.62
<i>Azospirillum</i> + PSB	1.22	5.40	102.67	184.81	165302	2.93
SEm±	0.02	0.11	1.94	4.45	5342	0.10
CD (p=0.05)	0.07	0.31	5.57	12.80	15354	0.28

low cost involvement in nutritional supply to the crop plants and higher bulb yield of onion obtained under this treatment. Similar results have been reported by Yetho *et al.*, (2012) and Patel *et al.*, (2011).

## References

- Arnon, D.I. 1949. Copper enzyme in isolated chloroplast. Polyphenol oxidase in *Beta vulgaris*. *Plant Physiology*, 24: 1-5.
- Aykrody, W.R. 1963. ICMR. Special Report, Series, No. 42.
- Barakart, M.A.S. and Gabr, S.M. 1998. Effect of different biofertilizer and nitrogen fertilizer levels on tomato plant. *Alexandria Journal Agril. Reseach.*, 43: 149-160.
- Bose, T.K., Kabir, J., Das, P. and Joy, R.P. 2001. *Organic farming in horticulture crops*. Tropical Horticulture, 2:20, Publishing Naya Prakash, New Delhi, p.150
- Choudhary, R.S., Das, Anchal, Patnaik, V.S. and Das, A. 2003. Organic farming for vegetable production using vermicompost and FYM in Kokriguda watershed of Orissa. *Indian Journal of Soil Science Conservation*, 31: 203-206.
- Fisher, R.A. 1950. Statistical method for research workers. Oliver and Boyd, Edinburg, London.
- Jackson, M.L. 1973. Soil chemical analysis. Prentice Hall of India Pvt.Ltd., New Delhi, pp.498.
- Mengistu F and Singh N, 1999. *Vegetable Science*, 26, pp.193-195.
- Mohandas S.1999. Biofertilizer for Horticulture crops. *Indian Horticulture*. 43: 32-37.
- Muthuramalingam S, Natarajan S, Sendurkumaran S and Muthuvel I. 2001. Morphological characters as influenced by spacing and nutrients in seed propagated aggregatum onion (*Allium cepa* L. var aggregatum Don.) type Gnanamedu local. *Madras Agricultural Journal*. 88: 379-382.
- Nadkarni KN.1954. *Allium cepa* Linn and *Allium sativum* Linn. *Indian Materia Medica*, Popular book depot, Bombay: 963.
- Paul NB, Rewari RB, Sen A, Sundeo WVB and Bhatnagar RS.1971. Rhizobial inoculation o pulse crops for better yield .Division of microbiology, New vistas in pulse production, IARI, New Delhi, pp-47-56.
- Ramesh PS, Singh M, Panwar NR, Singh AB and Ramana S. 2006. Response of pigeonpea varieties to organic manures and their influence of fertility and enzyme activity of soil. *Indian Journal of Agricultural Science*, 76: 252-254.
- Sharma RP, Dutt N and Chander G. 2009. Effect of vermicompost, farm yard manure and chemical fertilizers on yield, nutrient uptake and soil fertility in okra (*Abelmoschus esculentus*), onion (*Allium cepa*) sequence in wet temperature zone of Himachal Pradesh. *Journal of the Indian Society of Soil Science*, 57: 3.
- Snell PD and Snell GT. 1949. *Calorimetric methods of analysis*, 3<sup>rd</sup> End. Vol. II D. Van. Nostrand CO. Inc., New York.

- Tabatabai MA and Bremner JM. 1970. A simple turbidimetric method of determining total sulphur in plant material. *Agron. Journal*, 62: 805-806.
- Tanwar SPS, Sharma GL and Chahar MS. 2003. Effect of phosphorus and biofertilizer on yield , nutrient concentration and uptake by blackgram [*Vigna munga* (L.) Hepper]. *Legume Research*, 26: 39-41.
- Thanunathan K; Natarajan S; Kumar RS and Arul-Murugan K. 1997. Effect of different sources of organic amendments on growth and yield of onion in mine soil. *Madras Agriculture Journal*, 84: 382-384.
- Tilak KVBR and Annapurna K. 1993. Bacterial fertilizers. *Proceeding Indian National Academic Science*, 59 (3-4): 315-324.
- Yeptho KA, Vitoli AK, Singh SP, Kanaujia VB and Singh B. 2012. Quality production of *kharif* onion (*Allium cepa*) in response to bio-fertilisers inoculated organic manures. *Indian Journal of Agricultural Sciences*, 52: 2012.